



Goodwin, C., Berman, J., Janussen, D., Gocke, C., & Hendry, K. (2016). Hexactinellida (Porifera) from the Drake Passage (Southern Ocean) with a description of three new species. *Zootaxa*, 4126(2), 207-220. <https://doi.org/10.11646/zootaxa.4126.2.2>

Peer reviewed version

Link to published version (if available):
[10.11646/zootaxa.4126.2.2](https://doi.org/10.11646/zootaxa.4126.2.2)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Magnolia Press at <http://biotaxa.org/Zootaxa/article/view/zootaxa.4126.2.2>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Figure 1 – Sampling locations. A, Sars Seamount, B Shackleton Fracture Zone. Inset: box indicates the area shown in larger figure.




Figure 2. *Aulocalyx irregularis*, specimen BELUM.Mc2015.308. A. Collected specimen, scale in cm; B. Skeletal framework, Scale bar 1000 μ m, SEM; C. Pentactin, scale bar 100 μ m, SEM; D. Discohexaster, scale bar 10 μ m, SEM; E. Rhopalaster, scale bar 100 μ m, SEM.




Figure 3. *Doconesthes robinsoni* sp. nov. Type specimen MNHNCL POR-15002. A. Freshly collected specimen, scale in cm; B. Rehydrated specimen, scale in cm; C. Atrium of rehydrated specimen, scale in cm; D. Hypodermal pentactin, illustration traced from light microscope image, scale 100 μ m; E. Atrial hexactine, illustration traced from light microscope image, scale 100 μ m; F. End of choanosomal diactin, scale 200 μ m, light microscope; G. Dermal diactin, scale 100 μ m, light microscope; H. Hemioxyhexaster, scale 10 μ m, SEM; I. Strobiloplumicome, scale 10 μ m, SEM, the frontal ray is missing.




Figure 4. *Sympagella walleri* sp. nov. Type specimen MNHNCL POR-15003, strobiloplumicome image G from BELUM.Mc2015.313. A. Collected specimen; B. End of diactin, scale bar 200 μ m, light microscope; C. Dermal pentactin, scale bar 100 μ m, illustration traced from light microscope image; D. Dermal hexactin, scale bar 100 μ m, illustration traced from light microscope image; E. Pinular hexactin, scale bar 10 μ m, SEM; F. Discohexaster, scale bar 10 μ m, SEM; G, H. Strobiloplumicome, scale bar 10 μ m, SEM.




Figure 5. *Caulophacus palmeri* sp. nov. Type specimen MNHNCL POR_15001. A. collected specimen, scale in cm; B. end of diactin, scale bar 10 μ m, SEM; C. pentactin, scale bar 100 μ m, illustration traced from light microscope image; D. hexactin, scale bar 100 μ m, illustration traced from light microscope image; E. pinular hexactin, scale bar 10 μ m, SEM; F. discohexactin, scale bar 10 μ m, SEM.




Figure 6. *Rossella antarctica*: A. Collected specimen BELUM.Mc2015.284; B. Specimen BELUM.Mc2015.335; C. Rays of a calycocome from BELUM.Mc2015.284 (scale bar 10µm); D. Microdiscohexaster from BELUM.Mc2015.284 (scale bar 10 µm).

